



---

## Thermospheric Data Assimilation

**Tomoko Matsuo**

**REGENTS OF THE UNIVERSITY OF COLORADO THE**

---

**05/05/2016**

**Final Report**

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory  
AF Office Of Scientific Research (AFOSR)/ RTB1  
Arlington, Virginia 22203  
Air Force Materiel Command

**REPORT DOCUMENTATION PAGE**

 Form Approved  
 OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

**PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.**

1. REPORT DATE (DD-MM-YYYY) 30-04-2016			2. REPORT TYPE Final Report		3. DATES COVERED (From - To) from 01-Feb-13 to 31-Jan-16	
4. TITLE AND SUBTITLE Thermospheric Data Assimilation			5a. CONTRACT NUMBER			
			5b. GRANT NUMBER FA9550-13-1-0058			
			5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S) Dr. Tomoko Matsuo Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309-0216			5d. PROJECT NUMBER			
			5e. TASK NUMBER			
			5f. WORK UNIT NUMBER			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) REGENTS OF THE UNIVERSITY OF COLORADO THE OFFICE OF CONTRACTS AND GRANTS, 3100 MARINE ST RM 479, 572 UCB, BOULDER CO 80303-1058					8. PERFORMING ORGANIZATION REPORT NUMBER NA	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) USAF, AFRL DUNS 143574726 AF OFFICE OF SCIENTIFIC RESEARCH 875 NORTH RANDOLPH STREET, RM 3112 ARLINGTON VA 22203					10. SPONSOR/MONITOR'S ACRONYM(S) NA	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S) NA	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; Distribution is unlimited						
13. SUPPLEMENTARY NOTES Program Manager: Dr. Julie J Moses, AFOSR, 703-696-9586, Julie.moses@us.af.mil						
14. ABSTRACT This project demonstrates how the current limit of thermospheric mass density predictability can be extended by systematically integrating observations into a coupled thermosphere-ionosphere first-principles model. An ensemble data assimilation procedure, constructed with the NCAR Data Assimilation Research Testbed (DART) and the NCAR Thermosphere-Ionosphere Electrodynamics General Circulation Model (TIEGCM), can take advantage of the tight coupling between the ionosphere and thermosphere, enabling the inference of thermospheric temperature and compositions from abundant GPS-based ionospheric observations. Observing system simulation experiments suggest that thermospheric states, particularly temperature, can be well inferred by assimilating electron density profiles obtained from the COSMIC/FORMOSAT-3 mission into the TIEGCM. This in turn leads to a significant improvement of the neutral mass density forecasting longer than 3 days. Furthermore, validation of assimilation analyses with independent CHAMP mass density observations confirms that the approach indeed improves the thermospheric mass density specification. Predictability of the ionosphere can also be extended considerably by the approach developed in this project.						
15. SUBJECT TERMS Data assimilation, Ensemble forecasting, Thermosphere-ionosphere coupled data assimilation, Neutral mass density specification and forecasting,						
16. SECURITY CLASSIFICATION OF: a. REPORT U			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON Tomoko Matsuo 19b. TELEPHONE NUMBER (Include area code) 303-449-532	

## INSTRUCTIONS FOR COMPLETING SF 298

- 1. REPORT DATE.** Full publication date, including day, month, if available. Must cite at least the year and be Year 2000 compliant, e.g. 30-06-1998; xx-06-1998; xx-xx-1998.
- 2. REPORT TYPE.** State the type of report, such as final, technical, interim, memorandum, master's thesis, progress, quarterly, research, special, group study, etc.
- 3. DATES COVERED.** Indicate the time during which the work was performed and the report was written, e.g., Jun 1997 - Jun 1998; 1-10 Jun 1996; May - Nov 1998; Nov 1998.
- 4. TITLE.** Enter title and subtitle with volume number and part number, if applicable. On classified documents, enter the title classification in parentheses.
- 5a. CONTRACT NUMBER.** Enter all contract numbers as they appear in the report, e.g. F33615-86-C-5169.
- 5b. GRANT NUMBER.** Enter all grant numbers as they appear in the report, e.g. AFOSR-82-1234.
- 5c. PROGRAM ELEMENT NUMBER.** Enter all program element numbers as they appear in the report, e.g. 61101A.
- 5d. PROJECT NUMBER.** Enter all project numbers as they appear in the report, e.g. 1F665702D1257; ILIR.
- 5e. TASK NUMBER.** Enter all task numbers as they appear in the report, e.g. 05; RF0330201; T4112.
- 5f. WORK UNIT NUMBER.** Enter all work unit numbers as they appear in the report, e.g. 001; AFAPL30480105.
- 6. AUTHOR(S).** Enter name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. The form of entry is the last name, first name, middle initial, and additional qualifiers separated by commas, e.g. Smith, Richard, J, Jr.
- 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES).** Self-explanatory.
- 8. PERFORMING ORGANIZATION REPORT NUMBER.** Enter all unique alphanumeric report numbers assigned by the performing organization, e.g. BRL-1234; AFWL-TR-85-4017-Vol-21-PT-2.
- 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES).** Enter the name and address of the organization(s) financially responsible for and monitoring the work.
- 10. SPONSOR/MONITOR'S ACRONYM(S).** Enter, if available, e.g. BRL, ARDEC, NADC.
- 11. SPONSOR/MONITOR'S REPORT NUMBER(S).** Enter report number as assigned by the sponsoring/monitoring agency, if available, e.g. BRL-TR-829; -215.
- 12. DISTRIBUTION/AVAILABILITY STATEMENT.** Use agency-mandated availability statements to indicate the public availability or distribution limitations of the report. If additional limitations/ restrictions or special markings are indicated, follow agency authorization procedures, e.g. RD/FRD, PROPIN, ITAR, etc. Include copyright information.
- 13. SUPPLEMENTARY NOTES.** Enter information not included elsewhere such as: prepared in cooperation with; translation of; report supersedes; old edition number, etc.
- 14. ABSTRACT.** A brief (approximately 200 words) factual summary of the most significant information.
- 15. SUBJECT TERMS.** Key words or phrases identifying major concepts in the report.
- 16. SECURITY CLASSIFICATION.** Enter security classification in accordance with security classification regulations, e.g. U, C, S, etc. If this form contains classified information, stamp classification level on the top and bottom of this page.
- 17. LIMITATION OF ABSTRACT.** This block must be completed to assign a distribution limitation to the abstract. Enter UU (Unclassified Unlimited) or SAR (Same as Report). An entry in this block is necessary if the abstract is to be limited.

## **AWARD No FA9550-13-1-0058**

### **Thermospheric Data Assimilation**

**Principal Investigator: Dr. Tomoko Matsuo, University of Colorado at Boulder**

**Program Manager: Dr. Julie J. Moses, AFOSR/RTB1**

#### **Abstract**

Thermospheric mass density variation is the major source of aerodynamic drag estimation errors at altitudes below about 700 km. This project demonstrates how the current limit of thermospheric mass density predictability can be extended by systematically integrating observations into a coupled thermosphere-ionosphere first-principles model. An ensemble data assimilation procedure, constructed with the NCAR Data Assimilation Research Testbed (DART) and the NCAR Thermosphere-Ionosphere Electrodynamics General Circulation Model (TIEGCM), can take advantage of the tight coupling between the ionosphere and thermosphere, enabling the inference of thermospheric temperature and compositions from abundant GPS-based ionospheric observations. Observing system simulation experiments suggest that thermospheric states, particularly temperature, can be well inferred by assimilating electron density profiles obtained from the COSMIC/FORMOSAT-3 mission into the TIEGCM. This in turn leads to a significant improvement of the neutral mass density forecasting longer than 3 days. Furthermore, validation of assimilation analyses with independent CHAMP mass density observations confirms that the approach indeed improves the thermospheric mass density specification. Predictability of the ionosphere can be extended considerably by estimating neutral compositions from GPS-based ionospheric observations by the coupled thermosphere-ionosphere data assimilation approach developed in this project.

#### **Background**

Even though the Earth's upper atmosphere density is tenuous, it is substantial enough to exert significant drag on orbiting spacecraft and debris. Understanding this drag is complicated by the fact that this region is subject to highly variable external drivers from above and below. Also its ionized constituents affect telecommunication and navigation, motivating numerous observational and modeling efforts since the dawn of space exploration. Because of a lack of global observations of thermospheric parameters, our understanding of the density variability rests largely on the averaging of numerous observational samples, limiting our modeling capability to the realm of climatology. To overcome this shortage, the inference of thermospheric parameters from relatively plentiful ionospheric data is aided by exploiting our knowledge of the intimate coupling between the thermosphere and ionosphere using the ensemble Kalman filtering (EnKF) data assimilation system, constructed with the NCAR Data Assimilation Research


Testbed [Anderson et al., 2009] and the Thermosphere-Ionosphere-Electrodynamics General Circulation Model (TIEGCM) [Richmond et al., 1992]. The novelty of this approach [Matsuо et al., 2013] allows taking advantage of the expansive increase in recent years of the ability of GPS systems to globally monitor ionospheric parameters. Inferred thermospheric parameters are compared to independent observations obtained from the accelerometer on board the Challenging Minisatellite Payload (CHAMP) [Reigber et al., 2002].

## Finding 1

### **“Assimilation of COSMIC/FORMOSAT-3 electron density data into a coupled thermosphere-ionosphere general circulation model can extend the predictability of thermospheric mass density more than 3 days”**

The observing system simulation experiments (OSSEs) are conducted for a geomagnetically quiet 12-hour period from 0000 UT through 1200 UT of March 20, 2009, with a one-hour assimilation window and 90 ensemble members. Note that 90-member ensemble simulations are not centered at the truth from which synthetic observations for OSSEs are sampled, so the model ensemble is not entirely free of model biases. This 12-hour data assimilation period is followed by the 72-hour ensemble forecast period. Ensemble forecast simulations are initialized by the assimilation analyses obtained by the OSSEs. To examine the impact of data assimilation, additional ensemble simulations are run without data assimilation and referred to as the control. The experiment setup is described in detail in Hsu et al. [2016]. Figure 1 compares the neutral mass density errors from three 90-member ensemble forecast experiments, which are initialized with results from different OSSEs, to the control case. These OSSEs are executed with different filters constructed with different sets of the TIEGCM state variables being included in the ensemble Kalman filtering state vector. Suppose  $\mathbf{f}$  represents a vector of a given TIEGCM model state variable defined on the model grid. The EnKF state vector for each filter is composed of (a) the electron density  $\mathbf{f}_{e^-}$ , the atomic oxygen ion  $\mathbf{f}_{O^+}$ , and the temperature  $\mathbf{f}_T$ ; (b)  $\mathbf{f}_{e^-}$ ,  $\mathbf{f}_{O^+}$ ,  $\mathbf{f}_T$ , the atomic oxygen mixing ratio  $\mathbf{f}_O$ , and the molecular oxygen mixing ratio  $\mathbf{f}_{O_2}$ ; and (c)  $\mathbf{f}_{e^-}$ ,  $\mathbf{f}_{O^+}$ ,  $\mathbf{f}_T$ ,  $\mathbf{f}_O$ ,  $\mathbf{f}_{O_2}$ , as well as the horizontal neutral winds  $\mathbf{f}_u$ ,  $\mathbf{f}_v$ . Note that the sum of the mixing ratios of atomic oxygen, molecular oxygen, and molecular nitrogen is set to 1 in the TIEGCM, so the molecular nitrogen mixing ratio is adjusted according to the changes in the atomic oxygen mixing ratio  $\mathbf{f}_O$  and the molecular oxygen mixing ratio  $\mathbf{f}_{O_2}$ , even though it is not included in the state vector.

The comparison suggests that thermospheric states, particularly temperature, can be well inferred by assimilating COSMIC/FORMOSAT-3 data into the TIEGCM, which leads to an improvement of the neutral mass density forecast by more than 3 days. Effects of the neutral composition initialization, though smaller than the impact of neutral temperature initialization, are long-lasting when combined with the neutral wind initialization.




**Figure 1.** The root-mean-squared error (RMSE) of the neutral mass density over the course of the 72-hour ensemble forecast period. RMSE of the control experiment, in which no data is assimilated during the preceding 12-hour assimilation, is shown in gray. RMSE of the 90-member ensemble forecast experiment is shown in light green, yellow, and dark green, respectively, for model initialization with the three different filters described in the main text.

## Finding 2

### “Thermospheric mass density can be estimated from COSMIC/FORMOSAT-3 electron density data”

Actual COSMIC/FORMOSAT-3 electron density profile data are assimilated into the TIEGCM for a geomagnetically quiet period of 23-25 June, 2008 using the ensemble data assimilation experiment setup described in Lee et al. [2012]. In this experiment, the EnKF state vector is composed of  $\mathbf{f}_{e^-}$ ,  $\mathbf{f}_T$ ,  $\mathbf{f}_U$ ,  $\mathbf{f}_V$ ,  $\mathbf{f}_O$ ,  $\mathbf{f}_{O_2}$ . A one-hour assimilation window and 90 ensemble members are used. To validate the neutral mass density analysis resulting from this assimilation experiment, the result is compared to independent CHAMP in-situ mass density observations. During this period, the CHAMP satellite orbits at 330-360 km along the noon-midnight local time plane 30 times.



**Figure 2.** The root-mean-squared difference (RMSD) of CHAMP observations with the neutral mass density analyses and the control simulation is shown in red and blue, respectively, with the noon and midnight orbit displayed by solid and dashed line.

The squared difference between the CHAMP neutral mass density and the density analysis evaluated at the CHAMP location is computed and sorted for the noon and midnight orbits with respect to  $10^\circ$  latitude bins, and summarized as the RMSD as shown in Figure 3. For the noon orbit, the neutral mass density inferred from the assimilation of COSMIC/FORMOSAT-3 electron density observation agrees better with the CHAMP observations than the ensemble control simulations. For the midnight orbit, the neutral mass density specification is not improved by data assimilation over the control simulation.

### Finding 3

#### “Thermospheric composition can be estimated from GPS-based electron density data”

Predictability of the ionosphere can be extended considerably by estimating neutral compositions from GPS-based ionospheric observations by the coupled thermosphere-ionosphere data assimilation approach developed in this project. The fact that the ionospheric forecast is aided by the inference of thermospheric compositions is confirmed within the context of OSSEs for the COSMIC I mission [Hsu et al., 2014; 2016] and for the COSMIC II mission [Lee et al., 2013]. Furthermore, Chartier et al. [2016] and Chen et al. [2016] adopted the approach to assimilation of actual ground-based TEC data during storm periods, and showed that the approach indeed extended the

ionospheric forecast capability of a coupled thermosphere-ionosphere general circulation model. Though these are indirect evidences, thermospheric compositions can be estimated from both space-based and ground-based GPS observations to the extent that results in a considerable improvement of the ionospheric forecast by up to 24 hours.

### Peer-review Publications Supported by This AFOSR Award (8)

**Matsuо, T.**, I. T. Lee, and J. L. Anderson (2013), Thermospheric mass density specification using an ensemble Kalman filter, *J. Geophys. Res. Space Physics*, 118, 1339–1350, doi:10.1002/jgra.50162.

**Matsuо, T.** (2013), Upper atmosphere data assimilation with an ensemble Kalman filter, *AGU Monograph on modeling the Ionosphere-Thermosphere system* edited by J. D. Huba et al., ISBN: 978-0-87590-491-7.

Lee, I. T., H. F. Tsai, J. Y. Liu, C. H. Lin, **T. Matsuо**, and L. C. Chang (2013), Modeling impact of FORMOSAT-7/COSMIC-2 mission on ionospheric space weather monitoring, *J. Geophys. Res. Space Physics*, 118, 6518–6523, doi:10.1002/jgra.50538.

Hsu, C.-T., **T. Matsuо**, W. Wang, and J.-Y. Liu (2014), Effects of inferring unobserved thermospheric and ionospheric state variables by using an Ensemble Kalman Filter on global ionospheric specification and forecasting, *J. Geophys. Res. Space Physics*, 119, 9256–9267, doi:10.1002/2014JA020390.

Chartier, A. T., **T. Matsuо**, J. L. Anderson, N. Collins, T. J. Hoar, G. Lu, C. N. Mitchell, A. J. Coster, L. J. Paxton, and G. S. Bust (2016), Ionospheric data assimilation and forecasting during storms, *J. Geophys. Res. Space Physics*, 121, 764–778, doi:10.1002/2014JA020799.

Chen, C. H., W. H. Chen, C. H. Lin, I. T. Lee, **T. Matsuо**, J. T. Lin, J. Y. Liu, and C. T. Hsu (2016), Ionospheric data assimilation with TIE-GCM and GPS-TEC during geomagnetic storm conditions, *J. Geophys. Res. Space Physics*, under review.

**Matsuо, T.**, and C.-T. Hsu (2016), Ensemble data assimilation for thermospheric mass density specification and forecasting, *Progress in Earth and Planetary Science*, under review.

Hsu, C.-T., **T. Matsuо**, W. Wang, and J.-Y. Liu (2016), Seasonal and Solar activity dependence of a coupled thermosphere-ionosphere ensemble data assimilation system for ionospheric forecasting: FORMOSAT-3/COSMIC observing system simulation experiments, *Space Weather*, under review

## Other References

Anderson, J. L., T. Hoar, K. Raeder, H. Liu, N. Collins, R. Torn, and A. F. Arellano 397 (2009), The Data Assimilation Research Testbed: A community data assimilation 398 facility, *Bull. Am. Meteorol. Soc.*, 90, 1283–1296, 399 doi:10.1175/2009BAMS2618.1.

Lee, I. T., **T. Matsuo**, A. D. Richmond, J. Y. Liu, W. Wang, C. H. Lin, J. L. Anderson, and M. Q. Chen (2012), Assimilation of FORMOSAT-3/COSMIC electron density profiles into thermosphere/Ionosphere coupling model by using ensemble Kalman filter, *J. Geophys. Res. Space Physics*, 117, A10318, doi:10.1029/2012JA017700.

**Matsuo, T.**, and E. A. Araujo-Pradere (2011), Role of Thermosphere-Ionosphere Coupling in a Global Ionospheric Specification, *Radio Science*, 46, RS0D23, doi:10.1029/2010RS004576.

Reigber, C., H. Lühr, and P. Schwintzer (2002), CHAMP Mission Status, *Adv. Space Res.*, 30(2), 129-134, doi:10.1016/S0273-1177(02)00276-4.

Richmond, A. D., E. C. Ridley, and R. G. Roble (1992), A thermosphere/ionosphere general circulation model with coupled electrodynamics, *Geophys. Res. Lett.*, 19, 457 601-604.

1.

**1. Report Type**

Final Report

**Primary Contact E-mail**

**Contact email if there is a problem with the report.**

tomoko.matsuo@colorado.edu

**Primary Contact Phone Number**

**Contact phone number if there is a problem with the report**

3034490532

**Organization / Institution name**

University of Colorado at Boulder

**Grant/Contract Title**

**The full title of the funded effort.**

Thermospheric Data Assimilation

**Grant/Contract Number**

**AFOSR assigned control number. It must begin with "FA9550" or "F49620" or "FA2386".**

FA9550-13-1-0058

**Principal Investigator Name**

**The full name of the principal investigator on the grant or contract.**

Tomoko Matsuo

**Program Manager**

**The AFOSR Program Manager currently assigned to the award**

Julie J Moses

**Reporting Period Start Date**

02/01/2013

**Reporting Period End Date**

01/31/2016

**Abstract**

Thermospheric mass density variation is the major source of aerodynamic drag estimation errors at altitudes below about 700 km. This project demonstrates how the current limit of thermospheric mass density predictability can be extended by systematically integrating observations into a coupled thermosphere-ionosphere first-principles model. An ensemble data assimilation procedure, constructed with the NCAR Data Assimilation Research Testbed (DART) and the NCAR Thermosphere-Ionosphere Electrodynamics General Circulation Model (TIEGCM), can take advantage of the tight coupling between the ionosphere and thermosphere, enabling the inference of thermospheric temperature and compositions from abundant GPS-based ionospheric observations. Observing system simulation experiments suggest that thermospheric states, particularly temperature, can be well inferred by assimilating electron density profiles obtained from the COSMIC/FORMOSAT-3 mission into the TIEGCM. This in turn leads to a significant improvement of the neutral mass density forecasting longer than 3 days. Furthermore, validation of assimilation analyses with independent CHAMP mass density observations confirms that the approach indeed improves the thermospheric mass density specification. Predictability of the ionosphere can be extended considerably by estimating neutral compositions from GPS-based ionospheric observations by the coupled thermosphere-ionosphere data assimilation approach developed in this project.

**Distribution Statement**

DISTRIBUTION A: Distribution approved for public release.

This is block 12 on the SF298 form.

Distribution A - Approved for Public Release

**Explanation for Distribution Statement**

If this is not approved for public release, please provide a short explanation. E.g., contains proprietary information.

**SF298 Form**

Please attach your [SF298](#) form. A blank SF298 can be found [here](#). Please do not password protect or secure the PDF. The maximum file size for an SF298 is 50MB.

[AFD-070820-035\\_TM\\_Apr\\_2016.pdf](#)

**Upload the Report Document. File must be a PDF. Please do not password protect or secure the PDF . The maximum file size for the Report Document is 50MB.**

[FinalReport\\_ThermosphericDataAssimialtion\\_TM\\_Apr\\_2016.pdf](#)

**Upload a Report Document, if any. The maximum file size for the Report Document is 50MB.**

**Archival Publications (published) during reporting period:**

Matsuo, T., I. T. Lee, and J. L. Anderson (2013), Thermospheric mass density specification using an ensemble Kalman filter, *J. Geophys. Res. Space Physics*, 118, 1339–1350, doi:10.1002/jgra.50162.

Matsuo, T. (2013), Upper atmosphere data assimilation with an ensemble Kalman filter, AGU Monograph on modeling the Ionosphere Thermosphere system edited by J. D. Huba et al., ISBN: 978-0-87590 491-7.

Lee, I. T., H. F. Tsai, J. Y. Liu, C. H. Lin, T. Matsuo, and L. C. Chang (2013), Modeling impact of FORMOSAT-7/COSMIC-2 mission on ionospheric space weather monitoring, *J. Geophys. Res. Space Physics*, 118, 6518–6523, doi:10.1002/jgra.50538.

Hsu, C.-T., T. Matsuo, W. Wang, and J.-Y. Liu (2014), Effects of inferring unobserved thermospheric and ionospheric state variables by using an Ensemble Kalman Filter on global ionospheric specification and forecasting, *J. Geophys. Res. Space Physics*, 119, 9256–9267, doi:10.1002/2014JA020390.

Lee, I. T., H. F. Tsai, J. Y. Liu, C. H. Lin, T. Matsuo, and L. C. Chang (2013), Modeling impact of FORMOSAT-7/COSMIC-2 mission on ionospheric space weather monitoring, *J. Geophys. Res. Space Physics*, 118, 6518–6523, doi:10.1002/jgra.50538

Chartier, A. T., T. Matsuo, J. L. Anderson, N. Collins, T. J. Hoar, G. Lu, C. N. Mitchell, A. J. Coster, L. J. Paxton, and G. S. Bust (2016), Ionospheric data assimilation and forecasting during storms, *J. Geophys. Res. Space Physics*, 121, 764–778, doi:10.1002/2014JA020799.

**Changes in research objectives (if any):**

**Change in AFOSR Program Manager, if any:**

**Extensions granted or milestones slipped, if any:**

**AFOSR LRIR Number**

**LRIR Title**

**Reporting Period**

**Laboratory Task Manager**

**Program Officer**

**Research Objectives**

**Technical Summary**

**Funding Summary by Cost Category (by FY, \$K)**

	Starting FY	FY+1	FY+2
Salary			
Equipment/Facilities			
Supplies			
Total			

**Report Document**

**Report Document - Text Analysis**

**Report Document - Text Analysis**

**Appendix Documents**

**2. Thank You**

**E-mail user**

May 01, 2016 01:24:15 Success: Email Sent to: tomoko.matsuo@colorado.edu